



## The effect of 300 mW, 830 nm laser on chronic neck pain: A double-blind, randomized, placebo-controlled study

Roberta T. Chow<sup>a,\*</sup>, Gillian Z. Heller<sup>c</sup>, Les Barnsley<sup>b</sup>

<sup>a</sup> Castle Hill Medical Centre, 269-271 Old Northern Road, Castle Hill, NSW 2154, Australia

<sup>b</sup> Department of Statistics, Macquarie University, NSW 2109, Australia

<sup>c</sup> Faculty of Medicine, University of Sydney, Department of Rheumatology, Repatriation Hospital, Concord, NSW 2139, Australia

Received 16 June 2005; received in revised form 23 March 2006; accepted 17 May 2006

---

### Abstract

A randomized, double-blind, placebo-controlled study of low-level laser therapy (LLLT) in 90 subjects with chronic neck pain was conducted with the aim of determining the efficacy of 300 mW, 830 nm laser in the management of chronic neck pain. Subjects were randomized to receive a course of 14 treatments over 7 weeks with either active or sham laser to tender areas in the neck. The primary outcome measure was change in a 10 cm Visual Analogue Scale (VAS) for pain. Secondary outcome measures included Short-Form 36 Quality-of-Life questionnaire (SF-36), Northwick Park Neck Pain Questionnaire (NPNQ), Neck Pain and Disability Scale (NPAD), the McGill Pain Questionnaire (MPQ) and Self-Assessed Improvement (SAI) in pain measured by VAS. Measurements were taken at baseline, at the end of 7 weeks' treatment and 12 weeks from baseline. The mean VAS pain scores improved by 2.7 in the treated group and worsened by 0.3 in the control group (difference 3.0, 95% CI 3.8-2.1). Significant improvements were seen in the active group compared to placebo for SF-36-Physical Score (SF36 PCS), NPNQ, NPAD, MPQVAS and SAI. The results of the SF-36 - Mental Score (SF36 MCS) and other MPQ component scores (afferent and sensory) did not differ significantly between the two groups. Low-level laser therapy (LLLT), at the parameters used in this study, was efficacious in providing pain relief for patients with chronic neck pain over a period of 3 months.

© 2006 International Association for the Study of Pain. Published by Elsevier B.V. All rights reserved.

**Keywords:** Neck pain; GaAlAs; 830 nm; Low-level laser therapy (LLLT); RCT

---